

Research Article

Laparoscopic Cholecystectomy Effects on Renal Function Tests Among Patients Undergoing the Surgery in Ali Ibn Abi Taleb Hospital in Zahedan (South East of Iran) In 2018.

Ahmad Reza Shahraki^{1*}, Elham Shahraki², Mahdi Mohammadi³, Reza Abaee⁴, Elahe Shahraki⁵, Ali Reza Khazayi¹, Abbas Ali Niazi⁶

¹General surgeon, Assistant professor, Surgery Department, Zahedan medical faculty, Zahedan University of Medical Sciences, Zahedan, Iran.

²Associated professor, Nephrologist, Department of Internal Medicine, Ali Ibne Abi Taleb Hospital, Zahedan University of Medical Sciences, Zahedan, Iran.

³Associated professor, Biostatistics Department, Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran.

⁴MD, General Physician, Medical faculty, Tehran, Iran.

⁵MD, Faculty of Dentistry, Dentist, Zahedan University of Medical Sciences, Zahedan, Iran.

⁶Associated professor, Pathologist, Department of Pathology, School of Medicine, Cellular and Molecular Research Centre, Ali Ibne Abi Taleb Hospital, Zahedan University of Medical Sciences, Zahedan, Iran.

Received: 01 October, 2023 Accepted: 01 November, 2023 Published: 06 November 2023

Abstract:

Background: Laparoscopic Cholecystectomy, one of the most common elective surgical procedures, is a minimally invasive surgical procedure for patients with gallbladder diseases.

Objectives: This study aimed to evaluate the effect of laparoscopic cholecystectomy on Renal function Tests among patients undergoing the surgery in Ali Ibn Abi Taleb Hospital in Zahedan (South east of Iran).

Methods: Following a quasi-experimental design, 21 cases scheduled for laparoscopic cholecystectomy for whatever reasons at Ali Ibn Abi Taleb Hospital in Zahedan (Iran) were selected. To study the serum levels Renal function Tests, blood samples were collected at three stages, i.e., before the surgery (time point 1), 30 minutes after pumping the carbon dioxide gas into the abdomen (time point 2), and 30 minutes after removing the last port from the abdomen (time point 3). Data were analyzed using repeated measures.

Results: According to the findings, laparoscopic cholecystectomy surgery only had a significant effect on the BUN at the 99% confidence level. Also, BUN was increased and significantly differed at time points 2 (30 minutes after pumping) and 3 (30 minutes after removing the port) compared to time point 1 (30 minutes before the surgery).

Conclusions: Generally, it can be argued that despite observing some variations in the BUN test during the laparoscopic surgery, the Cr level did not change. Hence, it can be considered as a safe surgical intervention for renal functions.

Keywords: Laparoscopic Cholecystectomy, Renal function tests, Surgery, Laparoscopy.

Introduction:

The prevalence of gallstones has significantly increased, such that only in the US about one million cases are diagnosed each year [1]. This disease imposes a significant therapeutic-economic burden on the healthcare systems, even in western countries, to the extent that between 10% - 20% of people in the European and American societies are suffering from gallstones [2, 3] and refer to hospital emergency rooms with acute abdominal pains [4]. In most cases, gallstones symptoms are not significant. As about 25% - 50% of those suffering from

gallstone develop several complications, it is necessary to remove their gallbladder [5] using surgical techniques such as laparoscopic cholecystectomy, which is highly common. Laparoscopic cholecystectomy is one of the most common abdominal surgeries and the standard treatment for gallstones and cholecystitis [6]. In addition, it is a noninvasive technique [7] intended to reduce the potential damages of the disease [8]. Apart from several advantages, laparoscopic cholecystectomy is associated with better homeostasis compared to open surgeries [9]. Due to several reasons, including the need for a

small incision, low pain, short duration of hospitalization, early recovery, early start of eating food, and, in general, the quick return to daily activities, the laparoscopic cholecystectomy is the preferred option to treat most of the cases who suffer from the gallbladder [10, 11]. According to the currently available evidence, about 0.04 of cases who underwent the laparoscopic cholecystectomy develop complications [12] such as bile duct injury, acute biliary tract obstruction extrahepatic biliary duct rupture, falling gallstones into the abdominal cavity, and the subsequent formation of an abscess. Moreover, it also may cause some intraoperative complications, such as hemodynamic changes during blowing the gas into the peritoneum, including cardiac output, increased systemic vascular resistance, hypertension, heart rate changes, and reduced respiratory capacity [13]. In addition, it may cause some changes in the body, such as changes in acid-base, pulmonary status, cardiovascular system, and hemodynamic. There are reported that attributed changes in the liver function to impaired portal vein flow, decreased venous flow, and variations in intracranial pressure [14]. A dysfunction in the blood flow to the liver not only may result in liver dysfunction but also can disrupt the production of proteins made in hepatocytes, including coagulation factors (PT, PTT, and INR) [15]. Some empirical evidence reported PT and PTT levels following laparoscopic cholecystectomy [15, 16]; however, the evidence are still inconclusive over changes in the INR factor [16]. But Laparoscopy induced AKI is related with both hormonal stimuli for renal vasoconstriction and increased intra-abdominal pressure, causing hypoxemia and tubular renal injury. In conclusion, very few cases of laparoscopy induced AKI in young patients with no previous renal disease have been reported. Surgeons must consider this complication in the differential diagnosis of postoperative AKI [17]. As old patients are more prone to have pre-existing renal dysfunction, these patients are at increased risk of renal complications associated with laparoscopic surgery. In addition, living kidney donors are commonly operated via a laparoscopic approach. The effects of laparoscopy on renal haemodynamics and their potential consequences on kidney function should therefore be acknowledged by nephrologists [18]. This study did to show Laparoscopic Cholecystectomy are safe for kidneys yet.

3. Methods

3.1. Research Design:

The current study was carried out following a quasiexperimental design and using a convenience sampling method (for screening).

3.2. Statistical Society and Participants:

The statistical society of the present study included those scheduled for laparoscopic cholecystectomy, for whatever reasons, at Ali ibn Abi Taleb Hospital in Zahedan. A total of 21 cases were recruited.

3.3. Inclusion and Exclusion Criteria:

The inclusion criteria, which were determined based on previous studies, were as follows: all patients with symptomatic

gallstones who based on ultrasonography findings had stones in their gallbladder, being aged 18 to 75 years, a maximum body mass index (BMI) of 40, normal levels of BUN and Cr markers, and being ready to undergo the surgery[13]. The exclusion criteria were patients' intolerance to general anesthesia, having irrevocable coagulation disorders, suffering from metastatic lesions, and those whose surgery was changed from laparoscopic to open surgery during the surgery [19].

3.4. Experimental Design:

The present study was carried out in a way that, before the surgery, a venous catheter was considered for the patients, and their anesthesia was initiated with a dose of propofol (2 - 3 mg/kg) and fentanyl (0.5 g/kg) in the first 5 minutes, and it continued with 200 - 250 μ g/dL of propofol per minute. Then, the orotracheal intubation was performed. During the surgery, their applicable ringer's solution was 20 cc/kg. The surgery was conducted using the four-port technique, i.e., a 10-mm umbilical port, which was the first port that entered the abdomen with the aid of a port needle after pumping the gas into the abdomen, a 10- mm xiphoid port, a 5-mm mid-clavicular port at the Murphy point, and a 5-mm mid-clavicular port in the adjacency of umbilicus. During the surgery, the intra-abdominal pressure was maintained by a maximum of 15 mmHg of carbon dioxide, and the patients were under continuous cardiac and Capnography monitoring. The patients' blood samples were collected at three different time points, i.e., before the surgery (time point 1), 30 minutes after pumping the carbon dioxide gas into the abdomen (time point 2), and 30 minutes after removing the last port from the abdomen (time point 3). The blood samples taken in these time points were examined concerning daily BUN, Cr serum levels and then were transferred to a laboratory under a cold chain for further investigations [14]. It should be noted that the current study was carried out without a sexual preference (women), and steps mentioned in the anesthesia service reviews and the protocol of the American Society of Anesthesiologists were followed; hence, no preoperative preventive measure was taken. Furthermore, there were no indications for the Intraoperative injection of low-molecular-weight heparin (LMWH) and antibiotics.

3.5. Statistical Analysis:

Data analysis was performed using SPSS Ver.22 by repeated measures and Bonferroni follow-up test.

Results:

Before examining the research hypotheses, The Kolmogorov-Smirnov test was applied to test for a normal distribution, which indicated the normality of all variables, as the significance values of the normality tests were greater than 0.05. In addition, the assumption of equal variances was examined for each variable by Levene's test, which resulted in significant levels for all investigated variables. Hence, the assumption of equal variances was observed.




Figure1: Time- Changes of BUN and Cr on SD
($p < 0/01$)




Figure2: Changes of BUN and Cr on Mean
($p < 0/01$)

It shows that BUN can be affected by laparoscopic surgery and it is significant.

It shows that Cr can be affected by laparoscopic surgery but it is not significant.

Conclusion:

This study demonstrated that the Laparoscopic cholecystectomy surgery was only effective in changing (i.e., increasing) the BUN and did not have any significant effects on Cr. Bun changes can be affected by low protein diet, dehydration before or during surgery. Pneumoperitoneum during laparoscopy results in transient Oliguria and decreased glomerular filtration and renal blood flow. The presence of oliguria and elevated serum creatinine is suggestive of acute renal injury. Serum cystatin C has been described as a new marker for the detection of this type of injury. In this study, our aim was to compare the glomerular filtration rate estimated using cystatin C levels with the rate estimated using serum creatinine in patients with normal renal function who were undergoing laparoscopic surgery. [20]

Thus, except for the impacts of Pneumoperitoneum on portal vein flow due to liver functions and their related in inflammatory responses, some intervening factors, such as age, BMI, surgery duration, and surgical injuries, affect (either directly or indirectly) can change the BUN. Accordingly, patients with risk factors such as senescence, obesity, or a long wait for laparoscopic surgery, are likely to have significant coagulation activations. They are considered a high-risk group who are prone to the development of deep vein thrombosis following the surgery, which certainly causes a kind of Thromboprophylaxis. In this regard, laparoscopic surgery activates the coagulation and Bronchiolitis pathways and changes the height of the bTG and coagulation tests. It is necessary to mention some limitations and biases of our study. Firstly, not having the opportunity to assess the functions of the pancreas and other organs in the abdomen during the surgery, due to its high cost. Secondly, not having the chance to examine the functions of the organs after the surgery to evaluate postoperative changes. Thirdly, all participants of the present study were females who none of them was admitted to public hospitals.

But sometimes Laparoscopy induced AKI is related with both hormonal stimuli for renal vasoconstriction and increased intra-abdominal pressure, causing hypoxemia and tubular renal injury. In conclusion, very few cases of laparoscopy induced AKI in young patients with no previous renal disease have been reported. Surgeons must consider this complication in the differential diagnosis of postoperative AKI (19). As old patients are more prone to have pre-existing renal dysfunction, these patients are at increased risk of renal complications associated with laparoscopic surgery. In addition, living kidney donors are commonly operated via a laparoscopic approach. The effects of laparoscopy on renal hemodynamic and their potential consequences on kidney function should therefore be acknowledged by nephrologists. [18].As in Previous researches Laparoscopic cholecystectomy have no effects on coagulation and Liver function tests(21).It Shows nowadays this way is Safe and useful but more researches is needed.

Conflict of Interests:

The authors have no conflicts of interests to declare.

Declarations:

Ethical Approval and Consent to participate: The content of this manuscript is in accordance with the declaration of Helsinki for Ethics. No committee approval was required. Oral and written consent to participate was granted by the patient families.

Consent for publication: “Written informed consent was obtained from the patient's legal guardian for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.”

Availability of supporting data: It is available.

Competing interests:The author declares that they have no competing financial interests and nothing to disclose.

Funding: Zahedan University of Medical Sciences.

Authors' contributions:

Ahmad Reza Shahraki is the surgeon of patient and writes this paper.

The authors declare that they have no competing financial interests and nothing to disclose.

Ali Reza Khazy assists the surgery, Mehdi Mohammadi calculate statics, Elham Shahraki involved in patient examination and *Inclusion and Exclusion Criteria*, Elahe Shahraki and Reza Abaee participate in collecting data's and Abbas Ali Niazi did the Lab exams.

Acknowledgements

Generally, it can be argued that despite observing some variations in the BUN test during the laparoscopic surgery, the Cr level did not change. Hence, it can be considered as a safe surgical intervention for renal and liver functions.

References:

1. Mendez-Sanchez N, Chavez-Tapia NC, Motola-Kuba D, Sanchez-Lara K, Ponciano-Rodriguez G, Baptista H, et al. Metabolic syndrome as a risk factor for gallstone disease. *World J Gastroenterol*. 2005;**11**(11):1653–7. doi: 10.3748/wjg.v11.i11.1653. [PubMed: 15786544]. [PubMed Central: PMC4305948].
2. Petelin JB. Surgical management of common bile duct stones. *Gastrointest Endoscopy*. 2002;**56**(6B):S183–9. doi:10.1067/mge.2002.1290131.
3. Lammert F, Sauerbruch T. Mechanisms of disease: the genetic epidemiology of gallbladder stones. *Nat Clin Pract Gastroenterol Hepatol*. 2005;**2**(9):423–33. doi: 10.1038/ncpgasthep0257. [PubMed: 16265433].
4. Volzke H, Baumeister SE, Alte D, Hoffmann W, Schwahn C, Simon P, et al. Independent risk factors for gallstone formation in a region with high cholelithiasis prevalence. *Digestion*. 2005;**71**(2):97–105. doi:10.1159/000084525. [PubMed: 15775677].
5. Agrawal S, Jonnalagadda S. Gallstones, from gallbladder to gut. *Postgraduate Med*. 2015;**108**(3):143–53. doi: 10.3810/pgm.2000.09.1.1212.
6. Das NT, Deshpande C. Effects of Intraperitoneal Local Anaesthetics Bupivacaine and Ropivacaine versus Placebo on Postoperative Pain after Laparoscopic Cholecystectomy: A Randomised Double Blind Study. *J Clin Diagn Res*. 2017;**11**(7):UC08–12. doi:10.7860/JCDR/2017/26162.10188. [PubMed: 28893013]. [PubMed Central: PMC5583910].
7. Shiva A. The predictive factors for the success of laparoscopic cholecystectomy in patients with acute cholecystitis referring to the Qazvin province. *Ghazvin Univ Med Sci*. 2018.
8. Bessler M, Stevens PD, Milone L, Hogle NJ, Durak E, Fowler D. Multimedia.article: Transvaginal laparoscopic cholecystectomy: laparoscopically assisted. *Surg Endosc*. 2008;**22**(7):1715–6. doi: 10.1007/s00464-008-9779-1. [PubMed: 18322746].

9. Joris JL. Anesthesia for Laparoscopic Surgery. *Miller's Anesthesia*. 7th ed. Churchill livingstone: Elsevier; 2010. p. 2185–202. doi:10.1016/b978-0-443-06959-8.00068-6.
10. Townsend CMBR, Evers BM, Mattox K. *Sabiston Textbook of surgery*. Canada: Elsevier; 2010. p. 2185–6.
11. ChangWT, Lee KT, HuangMC, ChenJS, Chiang HC, KuoKK, et al. The impact of body mass index on laparoscopic cholecystectomy in Taiwan: an oriental experience. *J Hepatobiliary Pancreat Surg*. 2009;**16**(5):648–54. doi: 10.1007/s00534-009-0102-x. [PubMed: 19387531].
12. Picchio M, De Angelis F, Zazza S, Di Filippo A, Mancini R, Pattaro G, et al. Drain after elective laparoscopic cholecystectomy. A randomized multicentre controlled trial. *Surg Endosc*. 2012;**26**(10):2817–22. doi:10.1007/s00464-012-2252-1. [PubMed: 22538671].
13. Karaca O, Pinar HU, Turk E, Dogan R, Ahiskalioglu A, Solak SK. Effects of Single-Dose Preemptive Pregabalin and Intravenous Ibuprofen on Postoperative Opioid Consumption and Acute Pain after Laparoscopic Cholecystectomy. *J Invest Surg*. 2019;**32**(3):189–95. doi:10.1080/08941939.2017.1386738. [PubMed: 29157034].
14. Lai H, Mo X, Yang Y, Xiao J, He K, Chen J, et al. Association between duration of carbon dioxide pneumoperitoneum during laparoscopic abdominal surgery and hepatic injury: a meta-analysis. *PLoS One*. 2014;**9**(8). e104067. doi: 10.1371/journal.pone.0104067. [PubMed: 25110982]. [PubMed Central: PMC4128655].
15. Ntourakis D, Sergentanis TN, Georgiopoulos I, Papadopoulou E, Liasis L, Kritikos E, et al. Subclinical activation of coagulation and fibrinolysis in laparoscopic cholecystectomy: do risk factors exist? *Int J Surg*. 2011;**9**(5):374–7. doi: 10.1016/j.ijssu.2011.02.011. [PubMed: 21371576].
16. Demiryas S, Donmez T, Erdem VM, Erdem DA, Hatipoglu E, Ferahman S, et al. Comparison of the effects of spinal epidural and general anesthesia on coagulation and fibrinolysis in laparoscopic cholecystectomy: a randomized controlled trial: VSJ Competition, 2(nd) place. *Wideochir Inne Tech Maloinwazyjne*. 2017;**12**(3):330–40. doi: 10.5114/wiitm.2017.70249. [PubMed: 29062459]. [PubMedCentral: PMC5649509].
17. Ignacio Aguirre–Allende1,*, Lander Gallego–Otaegui2, Jose Luis Elosegui–Aguirrezabala1, Carlos Placer–Galán1, and Jose Maria Enriquez–Navascués1. Laparoscopy-induced severe renal failure after appendectomy. *Journal of Surgical Case Reports*, 2019;3, 1–3. doi: 10.1093/jscr/rjz079. Case Report.
18. Sophie de Seigneux1, Claude-Eric Klopfenstein2, Christophe Iselin3 and Pierre-Yves Martin1. **The risk of acute kidney injury following laparoscopic surgery in a chronic kidney disease patient**. *NDT Plus* (2011) 4: 339–341 doi: 10.1093/ndtplus/sfr071 Advance Access publication 17 July 2011 Teaching Point (Section Editor: A. Meyrier).

19. Papaziogas B, Koutelidakis I, Kabaroudis A, Galanis I, Paraskevas G, Vretzakis G, et al. Modifications of coagulation and fibrinolysis mechanism in laparoscopic vs. open cholecystectomy. *Hepatogastroenterology*. 2007;**54**(77):1335–8. [PubMed: 17708249].
20. Rodrigo Moreira e Lima,I* Lais Helena Camacho Navarro,I Giane Nakamura,II Daneshvari R. Solanki,III Yara Marcondes Machado Castiglia,I Pedro Tadeu Galvaõ Vianna,I Eliana Marisa GanemI. Serum cystatin C is a sensitive early marker for changes in the glomerular filtration rate in patients undergoing laparoscopic surgery.
21. Examining the Effect of Laparoscopic Cholecystectomy on Coagulation Tests among Patients Undergoing the Surgery in Ali Ibn Abi Taleb Hospital in Zahedan in 2018. Ahmad

Reza Shahraki 1, *, Ali Reza Khazayi 1, Mahdi Mohammadi 2, Elham Shahraki 3, Elahe Shahraki 4 and Abbas Ali Niaz. *Zahedan J Res Med Sci*. 2021 July; 23(3):e92588.

Copyright (c) 2023 The copyright to the submitted manuscript is held by the Author, who grants the Clinical Medicine and Health Research Journal a nonexclusive license to use, reproduce, and distribute the work, including for commercial purposes.



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)